



11 October 2022


Marine Ecosystem Modeling

Advanced Users

Olivier Aumont, IRD-IPSL/LOCEAN, Paris, France



Marine biogeochemistry: numerous processes to consider





A great number of processes at play which are all coupled:

- Biological: photosynthesis, respiration, trophic interactions, ...
- Chemical
- Physical: sedimentation, aggregation, mixing, transport, ...

...

Observing and (hopefully) understand



Modeling: why?

- Hypotheses testing

If we add, remove or change something, what happens?

- Quantitative dynamical framework

Are some datasets and/or parameter estimates consistent?

- Assessing some unknown rates/parameters

Based on observations, can we estimate some rates/fluxes/properties that are otherwise difficult to measure

- Prediction/forecasting

What the ocean will look like at some point in the future (or in the (far) past)?

- Design of an observing system or campaign

What is the best sampling strategy?

Overview of the presentation

- **Biogeochemical and ecosystem modeling: the different steps**

Constructing a model

- An historical perspective

When did it start and where we are

- Representing the physiology of the organisms

The different ways to model the living compartments in a model

- Modeling functional biodiversity

PFT models, trait-based models, gene-centric models, ...

- Final words

Mass conservation in a fluid

Relevant for nutrients and planktonic organisms


$$\frac{\partial C}{\partial t} = \underbrace{-C \nabla \cdot \vec{v}}_{\text{Advection}} + \underbrace{\frac{\partial}{\partial z} K_z \frac{\partial C}{\partial z} C + \nabla_h \cdot (K_h \nabla_h C)}_{\text{Diffusion (non resolved processes)}} + \underbrace{SMS}_{\text{Biogeochemical Sources and Sinks}}$$

- **Modeling ocean biogeochemistry and (planktonic) ecosystems requires an ocean circulation model**
- Any biases in the simulated ocean dynamics produce biases in marine biogeochemical and ecosystem models
- All the challenges related to dynamical modeling are pertinent for ocean biogeochemical modeling

Mesoscale/submesoscale, Mixing, overflows, boundary layers, ...

A first challenge: The computing cost

- Better modeling of the ocean circulation (and of the environment) generally requires to increase the spatial resolution
- Better modeling of the ecosystem and biogeochemical processes generally requires to increase the number of processes and prognostic variables (tracers)



Overview of the presentation

- Biogeochemical and ecosystem modeling: the different steps

Constructing a model

- **An historical perspective**

When did it start and where we are

- Representing the physiology of the organisms

The different ways to model the living compartments in a model

- Modeling functional biodiversity

PFT models, trait-based models, gene-centric models, ...


- Final words

Ecosystem/biogeochemical modeling: an historical perspective

- The pioneering studies : date back to the middle of the XXth century
- The first marine ecosystem model : Fleming (1939)

$$\frac{dP}{dt} = \mu P - g(t)$$

Simulation of a diatom bloom in the Channel




The Control of Diatom Populations by Grazing.*)

By



Richard H. Fleming,
Scripps Institution of Oceanography,
University of California, La Jolla, California.

- The first NPZ-type model coupling the dynamics of nutrients, phytoplankton and zooplankton: Steele (1974)



Ecosystem/biogeochemical modeling: Fasham et al. (1990)

- They defined the structure and the formulation of NPZD-type models on which most existing biogeochemical/ecosystem models currently rely



- All studies were restricted to 0-D or quasi 0-D frameworks

Spatially resolved biogeochemical models

Geochemical



Particles

Euphotic Layer (100-150m)




HAMOCC (1990)

→ A single equation for the whole « biology » :

$$\text{export production} = f(\text{Temp}) \cdot f(\text{light}) \cdot \text{PO}_4^2 / (\text{Ks} + \text{PO}_4)$$



→ The simplest models used in the global ocean models



How were they performing?

- They were extremely cheap but that was necessary considering the computing power available at that time
- Long-term simulations were feasible (paleo, future, steady-state)
- They were doing a decent job at reproducing the large-scale annual-mean patterns

Phosphate distribution in the Pacific ocean



- Current models perform better, but not by a lot and a large part of the improvement comes from a better representation of ocean dynamics

The first large-scale ecosystem (biogeochemical) models

Geochemical

NPZD

GLOBAL BIOGEOCHEMICAL CYCLES, VOL. 7, NO. 2, PAGES 417-450, JUNE 1993


**A SEASONAL THREE-DIMENSIONAL ECOSYSTEM
MODEL OF NITROGEN CYCLING IN THE NORTH
ATLANTIC EUPHOTIC ZONE**

J. L. Sarmiento,¹ R. D. Slater,¹ M. J. R. Fasham,² H. W.
Ducklow,³ J. R. Toggweiler,⁴ and G. T. Evans ⁵

GLOBAL BIOGEOCHEMICAL CYCLES, VOL. 10, NO. 4, PAGES 559-583, DECEMBER 1996


**Effects of plankton dynamics on seasonal carbon fluxes
in an ocean general circulation model**

Katharina D. Six and Ernst Maier-Reimer
Max-Planck-Institut für Meteorologie, Hamburg, Germany




An example

Satellite




HAMOCC3.1



Six and Maier-Reimer (1996)

Models have become more and more complex




- More complex does not necessarily imply more realistic! (Anderson, 2005; Friedrichs et al., 2007; Ward et al., 2013)
- A huge set of (often) badly constrained parameters

A second challenge: tuning/evaluating the models

- Hand tuning: the most common way


10, 20, 100x, ...



We learn a lot from the model dynamics (intuitive knowledge)



- Data assimilation approaches

Numerous difficulties



A second often hidden challenge

- Quite surprisingly, this step is often overlooked or not reported, despite it is a critical step



Overview of the presentation

- Biogeochemical and ecosystem modeling: the different steps

Constructing a model

- An historical perspective

When did it start and where we are

- **Representing the physiology of the organisms**


The different ways to model the living compartments in a model

- Modeling functional biodiversity

PFT models, trait-based models, gene-centric models, ...

- Final words

Modeling cell physiology




$$\frac{dP}{dt} = \mu_{max} * \frac{N}{K_N + N}$$

1940s

Monod/
Redfield

1 tracer/ pool


Modeling cell physiology



1940s

Monod/
Redfield

1 tracer/ pool



1960s


Droop/
Caperon
Internal stores

2/5 tracers/ pool

$$\frac{dQ_N}{dt} = V_{max} \frac{N}{K_N + N} - \mu Q_N$$

$$\frac{dP}{dt} = \mu_{max} * \left(\frac{1 - \frac{Q_{min}}{Q_N}}{1 - \frac{Q_{min}}{Q_{max}}} \right) P$$

Modeling cell physiology



1940s

Monod/
Redfield

1 tracer/ pool


1960s

Droop/
Caperon
Internal stores

2/5 tracers/ pool

Most biogeochemical/ecosystem models belong to one or the other of these classes


Modeling cell physiology



1940s

Monod/
Redfield


1 tracer/ pool



1960s

Droop/
Caperon
Internal stores

2/5 tracers/ pool




1970s

Shuter, Shuler
Macro-molecular
models

~10 tracers/ pool


Modeling cell physiology



1940s

Monod/
Redfield


1 tracer/ pool



1960s

Droop/
Caperon
Internal stores

2/5 tracers/ pool



1970s

Shuter, Shuler
Macro-molecular
models

~10 tracers/ pool


2000s

Metabolic
Reconstruction
FBA

~100-1000 tracers/ pool

- Very promising, for instance to evaluate the benefits and costs of metabolic pathways
- Identification of new metabolic pathways

Difficulties



Simulated effect of variations in ATPM on metabolism in a diatom

Ahmad et al. (2020)

- Not feasible considering the current computational constraints. Needs coarse-grained techniques
- This level of information is not available for most organisms
- FBA approaches assume steady-state or successive quasi steady-states (dFBA)
- Requires to specify an objective function to optimize which is not always easy

Overview of the presentation

- Biogeochemical and ecosystem modeling: the different steps

Constructing a model

- An historical perspective

When did it start and where we are

- Representing the physiology of the organisms

The different ways to model the living compartments in a model


- **Modeling functional biodiversity**

PFT models, trait-based models, gene-centric models, ...


- Final words

Modeling (functional) biodiversity


- A great challenge for biogeochemical/ecosystem modeling is to represent biodiversity
- Biodiversity = functional biodiversity
- Biodiversity has important consequences on biogeochemistry and ecosystem functioning



Hillebrand et al. (2013)



Plankton Functional Types Models



Le Quéré et al. (2005)

- **How to define PFTs?**

A PFT should have a specific biogeochemical/ecosystem function

A PFT could be allocated a distinct set of physiological /environmental/food/... characteristics

A PFT should have some importance in a region of the ocean

- **Difficulties**

Numerous species are lumped into a limited number of boxes

How many boxes should be set? How to set a single set of parameters?

Fixed, a priori-defined structure of the model

Tuning becomes quickly a pain!


Trait-based models

- What is it?

Species are not specifically modeled

Organisms are identified by a few taxa-transcending properties: their key traits and the trade-offs between them

Structure and function of ecological communities emerge from properties of the individual organisms.



- Difficulties


What traits should be represented?

Quantifying the trade-offs is very often challenging. Metabolic reconstruction can be very promising.


All traits are (most of the time) accessible everywhere/all the time. Evolution?

Can be very very expensive (1 trait = 1 additional dimension to the problem)


Size is a master trait



- Many metabolic rates show some dependency to size (allometry) $Y = Y_0 W^b$
- Many processes/fluxes are impacted by size: sinking of particles, feeding strategy, motility, vertical migrations, ...
- Trophic interactions are influenced by size (who's eating whom?)
- Biomass distribution as a function of size often shows some regular properties (Sheldon et al., 1972)





Wirtz (2012)




Xu et al. (2021)

Most current models mix both approaches


- The different modeling approaches are not hermetically separated
- Most models mix to some extent PFT and trait-based modeling formalisms




(b) Size classes



(c) Functional groups



(d) Thermal norms




A version of the DARWIN model (Dutkiewicz et al., 2020)

Gene-centric models


- Bringing together (meta)genomic data and biogeochemical models is challenging
- A major difficulty is that they differ in the considered currency: 'omics data refer to genomes, proteins and metabolites; biogeochemical models refer to concentrations, biomass and biogeochemical functions
- A functional gene-centric approach: organisms are grouped according to their functional genes/metabolisms (Reed et al., 2014)
- As most organisms in the sea are uncultured, simulating their genes is impossible. An alternative is to randomly allocate genes from a know pool to construct a set of organisms (Cole et al., 2017). And the environment selects.

GENOME model

- A example of a gene-centric modeling study in the Atlantic Ocean (Cole et al., 2017)



Surface genes concentration in June



Metagenomes in the Amazon river plume in June clustered by similarities

- These approaches linking omics and biogeochemical/ecosystem are still in their infancy but are rapidly growing

Final words

- A brief and subjective overview of biogeochemical/ecosystem modeling
- Many aspects have been omitted: upper trophic levels, evolution, niche-modeling, micro-nutrients, diagenetic/benthic, ...
- Many challenges have not been mentioned (and I certainly do not know all of them)
- One of these (not clearly stated) challenges is to bring together an increasing number of very diverse expertise: mathematics, computer science, physics, physiology, biogeochemistry, ecology, 'omics, ...
- Models are not the real world. They are always imperfect and necessarily show some level of deficiency

PISCES

Basic information on the model options


Objectives of that presentation

- Not a description of the PISCES model. This is a session for advanced PISCES users!
- Not an exhaustive description of all PISCES secrets
- A brief description of the PISCES optional features that can be activated from the namelist
- It also describes some key parameters that modify the behavior of these features
- This is probably imperfect. Your inputs are welcome to improve that document
- A technical documentation of PISCES is still missing and should come (soon we hope)

PISCES-std vs. PISCES-QUOTA

`ln_p4z = .true.`


PISCES-std (24/25 tracers)




PISCES-std vs. PISCES-QUOTA

`ln_p5z = .true.`


PISCES-QUOTA (39/40 tracers)



Code structure - Main



Code structure - SMS




PISCES-std vs. PISCES-QUOTA (2)

- Most of the optional features work in these two main versions of PISCES
- The sediment module cannot be activated with PISCES-QUOTA (no variable stoichiometry in the sediment module)
- PISCES-QUOTA is significantly more expensive than PISCES-std (>2x)
- Many parameterization choices are common to both versions
- The rest of that presentation will be based on PISCES-std

Prognostic ligands

- In the default configuration, concentration of iron ligands is either :
 - 1) set to a constant value defined in the namelist (`ln_ligand`)
 - 2) or to a variable field diagnosed from DOC (`ln_ligvar` = .true.)
- A prognostic description of the ligands can be activated by setting
`ln_ligand = .true.`
- This adds a new prognostic tracer `jplgw` (25 prognostic tracers)
- `p4zligand` is now called which computes the sinks (rem mineralization, photodegradation)
- Various additional routines have some new codes activated (`p4zprod`, `p4zfechem`, ...)

Prognostic ligands (2)



Sediment model

In the default configuration, exchanges with the sediments are modeled based on a simple metamodel proposed by Middelburg et al. (1996):

$$F_{sed} = F(\text{NO}_3, \text{O}_2, Z, \dots)$$

A full prognostic diagenetic model is embedded in PISCES and can be activated by setting `ln_sediment = .true.`


This sediment model can be used in a standalone mode (without PISCES running) but the code needs to be compiled with the CPP key `key_sed_off`

When running with PISCES, the exchanges between the water column and the sediments can be 1-way or 2-ways: `ln_sed_2way`

A session is dedicated to this sediment model

Reactivity-continuum for POC

This parameterization is described in Aumont et al. (2017)





$$\bar{k} = \frac{\int_0^{\infty} k^{\nu} e^{-ak} e^{-kt} dk}{\int_0^{\infty} k^{\nu-1} e^{-ak} e^{-kt} dk} = \frac{\nu}{(a+t)}$$

No switch to activate that parameterization. The number of lability classes is set in the namelist by jcpoc (jcpoc = 1 is equivalent to no variable lability)

The shape of the gamma function controlling the initial distribution is set by rshape

Reactivity-continuum for POC (2)



	noRC	RC
n	2656	2656
r	0.7	0.77
RMSE	0.14	0.08
Bias	0.12	0.02
RI	4.9	1.6
MEF	-0.6	0.71

This parameterization is coded in p4zpoc

The computing cost is increased by ~10% for 15 lability classes


Diurnal vertical migration of mesozooplankton

Not a prognostic parameterization ! See Gorgues et al. (2019)

DVM parameterization is activated by `ln_dvm_meso = .true.`

Migration depth is parameterized according to Bianchi et al. (2013)

$$Z_{\text{mig}} = F(\text{O}_2, \text{Chl}, T)$$



From Bianchi et al., 2013

Diurnal vertical migration of mesozooplankton (2)

A constant fraction of mesozoo is prescribed to migrate ($x_{fracmig}$).

Microzoo is not migrating

Organisms are assumed to be at the surface at night and at the migration depth during daytime

Organisms are supposed to respire, excrete DOM and inorganic nutrients and egest fecal pellets in both habitats (function of daylength and temperature)

This parameterization is coded in p4zmeso

The computing cost is only modestly increased

Concluding remarks

This was a short description of the optional features that come with the standard version of PISCES

Several aspects that can be controlled from the namelist have not been mentioned (external inputs, grazing param., ...)

In addition to the standard version of the model, three alternative versions do exist

- 1) PISCES-ISO: includes a description of ^{15}N and ^{13}C
- 2) PISCES-GAS: DMS and N_2O are explicitly modeled
- 3) PISCES-BYONIC: Mn, Co, Zn cycles are represented in that version

They are or will be made available on a GITLAB server

Code structure

